

Clinical Frailty Scale Table

(ref 7)

1	Very fit	People who are robust, active, energetic and motivated. These people commonly exercise regularly. They are among the fittest for their age
2	Well	People who have no active disease symptoms but are less fit than category 1. Often, they exercise or are very active occasionally, e.g. seasonally
3	Managing well	People whose medical problems are well controlled, but are not regularly active beyond routine walking
4	Vulnerable	While not dependent on others for daily help, often symptoms limit activities. A common complaint is being “slowed up”, and/or being tired during the day
5	Mildly frail	These people often have more evident slowing, and need help in high order IADLs (finances, transportation, heavy housework, medications). Typically, mild frailty progressively impairs shopping and walking outside alone, meal preparation and housework
6	Moderately frail	People need help with all outside activities and with keeping house. Inside, they often have problems with stairs and need help with bathing and might need minimal assistance (cuing, standby) with dressing
7	Severely frail	Completely dependent for personal care, from whatever cause (physical or cognitive). Even so, they seem stable and not at high risk of dying (within ~ 6 months)
8	Very severely frail	Completely dependent, approaching the end of life. Typically, they could not recover even from a minor illness
9	Terminally ill	Approaching the end of life. This category applies to people with a life expectancy

- ❖ Clinical frailty score 1-4 can be triaged as level of priority 1 for physical fitness assessment in ICU priority level.
- ❖ Clinical frailty score 5-6 can be triaged as level of priority 2 for physical fitness assessment in ICU priority level.
- ❖ Clinical frailty score 7-9 can be triaged as level of priority 3 for physical fitness assessment in ICU priority level.

